

APPROVED	BY	CLASS	SUBCLASS

Applicant : Jonathan M. Friedman
Application No. : 0,592 Filed: July 20, 2001

Docket No.: FAZIX/001 US

Confirmation No.: 1403

For: A method for ab initio determination of macromolecular crystallographic phases at moderate resolution by symmetry enforced orthogonal multicenter spherical harmonic-spherical Bessel expansion.

Attorney: James F. Haley, Jr., Reg. No. 27,794 Tel: (212) 596-9000

Sheet 1 of 7

1/7



Analysis of Packing Function Solutions for Monomeric Proteins




FIG. 1

RECEIVED

APR 24 2003

TECH CENTER 1600/2900

APPROVED	C. FIG.	Applicant : Jonathan M. Friedman
BY	CLASS	Application No. : 0,592 Filed: July 20, 2001
CHAFFSMAN	SUBCLASS	For : A method for ab initio determination of macromolecular crystallographic phases at moderate resolution by symmetry enforced orthogonal multicenter spherical harmonic-spherical Bessel expansion.
		Attorney : James F. Haley, Jr., Reg. No. 27,794 Tel: (212) 596-9000

Docket No. : FAZIX/001 US
 Confirmation No. : 1403
 Sheet 2 of 7

2/7



Analysis of Packing Function Solutions for Monomeric Proteins




FIG. 2

RECEIVED

APR 24 2003

TECH CENTER 1600/2900

APPROVED	FIG.
BY	CLASS
DRAFTSMAN	SUBCLASS


Applicant : Jonathan M. Friedman
 Application No. : 0,592 Filed: July 20, 2001
 For : A method for ab initio determination of macromolecular crystallographic phases at moderate resolution by symmetry enforced orthogonal multicenter spherical harmonic-spherical Bessel expansion.
 Attorney : James F. Haley, Jr., Reg. No. 27,794 Tel: (212) 596-9000

Docket No. : FAZIX/001 US
 Confirmation No. : 1403
 Sheet 3 of 7




3/7

START:




For Use with a Radial Correction r with Modes 5 thru 7:



Calculation Mode-Specific Routines:

Modes 1 & 2
 (Unphased Diffraction Amplitudes to Phased FT of SHSB-modeled Unit Cell)



Modes 1 & 2 (cont'd)




FIG. 3

RECEIVED

APR 24 2003

TECH CENTER 1600/2900

APPROVED	CLASS	Fi
BY		
DRAFTSMAN		

Applicant : Jonathan M. Friedman
 Application No. : 09/910,592 Filed July 20, 2001

Docket No.: FAZIX/001 US
 Confirmation No. : 1403
 For a method for ab initio determination of macromolecular crystallographic phases at moderate resolution by symmetry enforced orthogonal multicenter spherical harmonic-spherical Bessel expansion.
 Attorney: James F. Haley, Jr., Reg. No. 27,794 Tel: (212) 596-9000 Sheet 4 of 7

4 / 7




Figure . A schematic example: Two choices for filling the same portion of a crystal unit cell from an orthorhombic Spacegroup. Although the spheres on the right are smaller than those on the left, for some crystals the local maximum in the packing on the right would be the packing of maximal consistency with the crystallographic data.

FIG. 4

RECEIVED

APR 24 2003

TECH CENTER 1600/2900

APPROVED	C. FIG.
BY	CLASS
DATE	SUBCLASS

Applicant : Jonathan M. Friedman
 Application No. : 0,592 Filed: July 20, 2001
 For: A method for ab initio determination of macromolecular crystallographic phases at moderate resolution by symmetry enforced orthogonal multicenter spherical harmonic-spherical Bessel expansion.
 Attorney : James F. Haley, Jr., Reg. No. 27,794 Tel: (212) 596-9000

Docket No. : FAZIX/001 US
 Confirmation No. : 1403
 Sheet 5 of 7



5/7




FIG. 5

RECEIVED

APR 24 2003

TECH CENTER 1600/2900



Identical Image from Expansions about Different Origins:

$$\begin{aligned}
 & 11.22 x + 0.93e^{2\pi i(1.0) x} + 4.31e^{2\pi i(0.4) x} + \dots = \text{Image 1} \\
 & 10.01 x + 0.31e^{2\pi i(0.78) x} + 8.31e^{2\pi i(1.2) x} + \dots = \text{Image 2}
 \end{aligned}$$

Image 1 and Image 2 are identical, as indicated by the bracket connecting the two results.

Symmetry Expanded Direct Space Basis Functions:

$$? x + ? e^{2\pi i(?) x} + ? e^{2\pi i(?) x} + \dots = \text{Expanded Image}$$

The diagram shows three rectangular boxes, each containing a different arrangement of spheres, representing the basis functions. These are summed to form a final rectangular box containing a complex, expanded image of the spheres.

With a properly chosen origin, 45-55% of the unit cell can be expanded. (Most protein crystals are > 45% solvent.)

FIG. 6

RECEIVED

APR 24 2003

TECH CENTER 1600/2900


APPROVED	FIG	Applicant
BY	CLASS	Application No. : 09/910,591 Filed: July 20, 2001
CRAFTSMAN		For : A method for ab initio determination of macromolecular crystallographic phases at moderate resolution-by-symmetry enforced orthogonal multicenter spherical harmonic-spherical Bessel expansion.

Docket No.: FAZIX/001
 Confirmation No. : 1403
 Attorney : James F. Haley, Jr., Reg. No. 27,794 Tel: (212) 596-9000 Sheet 7 of 7

7 / 7



Component Direct Space Basis Functions:



Component Fourier Transforms:

$$a_{001} F_{\text{solo}}^{001}(\text{hkl}) + a_{211} F_{\text{solo}}^{211}(\text{hkl}) + a_{111} F_{\text{solo}}^{111}(\text{hkl}) + \dots = F_{\text{obs}}(\text{hkl})$$

$$a_{001} = \sum_{\text{hkl}} F_{\text{solo}}^{*001}(\text{hkl}) F_{\text{obs}}(\text{hkl}) \quad [\text{presume } \phi = 0.00 \text{ to start}]$$

$$F_{\text{accum}}(\text{hkl}) = a_{001} F_{\text{solo}}^{001}(\text{hkl})$$

$$a_{211} = \sum_{\text{hkl}} F_{\text{solo}}^{*211}(\text{hkl}) (|F_{\text{obs}}(\text{hkl})| - |F_{\text{accum}}^n(\text{hkl})|) e^{2\pi i \phi_{\text{accum}}^n(\text{hkl})}$$

$$F_{\text{accum}}^{n+1}(\text{hkl}) = F_{\text{accum}}^n(\text{hkl}) + a_{211} F_{\text{solo}}^{211}(\text{hkl})$$

FIG. 7

RECEIVED

APR 24 2003

TECH CENTER 1600/2900